

# Case-based Adaptation Problems in Medicine

Rainer Schmidt, Olga Vorobieva, Lothar Gierl

Institut für Medizinische Informatik und Biometrie, Universität Rostock  
Rembrandtstr. 16 / 17, D-18055 Rostock, Germany  
[rainer.schmidt / olga.vorobieva / lothar.gierl] @medizin.uni-rostock.de

**Abstract.** So far, Case-based Reasoning has not become as successful in medicine as in some other application domains. One, probably the main reason is that on the one side in Case-based Reasoning the adaptation problem could not be solved in a general way. On the other side in medicine adaptation is often more difficult than in other domains, because usually more and complex features have to be considered. In this paper, we not only summarise experiences with adaptation in medicine, but we want to elaborate typical medical adaptation problems and hope to indicate possibilities how to solve them.

## 1 Introduction

Case-based Reasoning (CBR) has become a successful technique for knowledge-based systems in many domains, while in medicine some more problems arise to use this method. A CBR system has to solve two main tasks [AP94]: The first one is the retrieval, which is the search for or the calculation of most similar cases. For this task much research has been undertaken. The basic retrieval algorithms for indexing [SHK89], Nearest Neighbor match [Br90], pre-classification [Qu93] etc. have already been developed some years ago and have been improved in the recent years. So, actually it has become correspondingly easy to find sophisticated CBR retrieval algorithms adequate for nearly every sort of application problem.

The second task, the adaptation means modifying solutions of former similar cases to fit for a current problem. If there are no important differences between a current and a similar case, a solution transfer is sufficient. Sometimes only few substitutions are required, but usually adaptation is a complicated process. While in the early nineties the focus of the CBR community lay on retrieval, in the late nineties CBR researchers investigated various aspects of adaptation. Theories and models for adaptation [e.g. BW98, FM99] and even algorithms to consider the effort of adaptation already during the retrieval process [SK98] have been developed.

However, adaptation is still domain dependent. Usually, for each application specific adaptation rules have to be generated. Unfortunately, attempts to learn adaptation rules from case bases automatically [HK96] lead to an intractable amount of specific rules.

Since adaptation is even more difficult in medicine, we want to elaborate typical medical adaptation problems and we hope to show possibilities how to solve them.

## 2 Medical Case-based Reasoning systems

Though CBR was not as successful in medicine as in some other domains so far, several medical systems have already been developed which at least apply parts of the Case-based Reasoning method. Here we do not want to give a review of these systems (for that see [GBS98, Sc01]), but we intend to show the main developments concerning adaptation in this area.

### 2.1 Avoiding the adaptation problem

Some systems avoid the adaptation problem, because they do not apply the complete CBR method, but only a part of it, namely the retrieval. These systems can be divided into two groups, retrieval-only systems and multi modal reasoning systems.

Retrieval-only systems are mainly used for image interpretation. First, Macura [MM95] claimed that users of image interpretation systems themselves wanted to be responsible for adaptation. Later on Perner [Pe01] pointed out that image interpretation mainly is a classification task, which in more sophisticated systems is enriched by retrieval. However, retrieval-only systems are not only used for image interpretation, but for other visualisation tasks too, e.g. for the development of kidney function courses [Sc99] and for hepatic surgery [Du97].

Multi modal reasoning systems apply parts of different reasoning methods. From CBR they usually incorporate the retrieval step, often to calculate or support evidences [Mo00], e.g. in CARE-PARTNER [Bi98] CBR retrieval is used to search for similar cases to support evidences for a rule-based program.

### 2.2 Solving the adaptation problem

So far, only a few medical systems have been developed that apply the complete CBR method. In these systems mainly three techniques are used for adaptation: adaptation operators or rules, constraints, and Compositional Adaptation. Furthermore, abstracting from specific single cases to more general prototypical cases sometimes supports the adaptation.

**Adaptation rules and operators.** One of the earliest medical expert systems that use CBR techniques is CASEY [Ko88]. It deals with heart failure diagnosis. The most interesting aspect of CASEY is the ambitious attempt to solve the adaptation task. Since the creation of a complete rule base for adaptation was too time consuming, a few general operators are used for adaptation. Since many features have to be considered in the heart failure domain and since consequently many differences between cases can occur, not all differences between former similar cases and a current case can be handled by the developed general adaptation operators. So, if no similar case can be found or if adaptation fails, CASEY uses a rule-based domain theory.

However, the development of complete adaptation rule bases never became a successful technique to solve the adaptation problem in medical CBR systems, because the bottleneck for rule-based medical expert systems, the knowledge acquisition, not only occurs again, but also seems to be more difficult to overcome.

**Constraints.** A more successful technique is the application of constraints. In ICONS, an antibiotics therapy adviser [SG01], a similar case is retrieved, the list of

therapy recommendations made for the retrieved case is transferred to the current patient, and this list is reduced by additional contraindications of the current patient. In ICONS the adaptation is rather easy, it is just a reduction of a list of solutions (therapies) by constraints (contraindications).

Another example for applying constraints is a diagnostic program concerning dysmorphic syndromes. This program has been developed in the GS.52 project [GS94]. The retrieval provides a list of prototypes sorted according to their similarity in respect to the current patient. Each prototype defines a diagnosis (dysmorphic syndrome) and represents the typical features for this diagnosis. The provided list of prototypes is checked by a set of explicit constraints. These constraints state that some features of the patient either contradict or support specific prototypes (diagnoses). So the provided list of prototypes is reduced by contradictions and sorted anew concerning evidences.

A typical task for applying constraints is menu planing [PMS98], where different requirements have to be served: special diets and individual factors, not only personal preferences, but also contraindications and demands based on various complications.

**Compositional Adaptation.** A further successful adaptation technique is Compositional Adaptation [WSC98]. In TA3-IVF [Ju98], a system to modify in vitro fertilisation treatment plans, relevant similar cases are retrieved (the relevance has to be specified by the user) and Compositional Adaptation computes weighted averages for the solution attributes. The weights per retrieved case are determined by the similarity to the current patient; the values of the solution attributes are nominal valued.

In TeCoMed [SG02], an early warning system concerning threatening influenza waves, explicit similarity criteria guarantee that a list of the most similar former courses is retrieved. Compositional Adaptation is applied to this list. That means, the list is split into two lists concerning the question whether a warning was appropriate. For both new lists the sum of the similarities of their cases is computed and the comparison of both sums decides whether a warning should be issued.

**Abstraction.** As one reason for the adaptation problem is the extreme specificity of single cases, the generalisation from single cases into abstracted prototypes [GS94] or classes [Bi96] may support the adaptation. The idea of generating more abstract cases is typical for the medical domain, because (proto-) typical cases directly correspond to (proto-) typical diagnoses or therapies. In GS.52 [GS94] each case is characterised by a list of features, which usually contains between 40 and 130 symptoms and syndromes. This means, there are so many differences between a current and a similar case that adaptation cannot take all of them into account. So, for all cases with the same dysmorphic syndrome a prototype is created, which contains the most frequent observed features of these cases. Such an abstracted prototypical case represents a dysmorphic syndrome and usually contains only up to 20 features. For a current case the most similar prototypes are calculated. Subsequently, for adaptation only few constraints have to be checked.

While in GS.52 all prototypes are organised on the same level, in MNAOMIA [Bi96], a hierarchy of classes, cases, and concepts with few layers is used. MEDIC [Tu88], a diagnostic reasoner on the domain of pulmonology, consists of a multi-layered hierarchy of schemata, of scenes, and of memory organisation packets of individual cases.

Apart from the three above mentioned adaptation techniques, only few different attempts to solve the adaptation task in medicine have been undertaken. And even they somehow apply one of the three techniques, e.g. in CASIMIR [LB00], a program for breast cancer treatment, similarity paths and reformulation are used to support the adaptation, but adaptation knowledge in form of rules is still required.

### **3 Adaptation problems in endocrinology diseases therapy support**

Very recently we have developed some programs for endocrinology support in a children's hospital. Here, we do not present these programs in detail, but we focus on adaptation problems within them to illustrate general adaptation problems in medicine. These programs deal with therapy support, especially for hypothyroidism.

All body functions are regulated by the endocrine system. The endocrine gland produces hormones and secretes them in blood. Hypothyroidism means that a patient's thyroid gland does not produce enough thyroid hormone naturally. If hypothyroidism is undertreated, it may lead to obesity, bradycardia and other heart diseases, memory loss and many other diseases [Ha00]. Furthermore in children it causes mental and physical retard. If hypothyroidism is of autoimmune nature, it sometimes occurs in combination with further autoimmune diseases such as diabetes. The diagnosis hypothyroidism can be established by blood tests. The therapy is inevitable: thyroid hormone replacement by levothyroxine. The problem is to determine the therapeutic dose, because the thyroxin demand of a patient follows only very roughly general schema and so the therapy must be individualised [Gr94]. If the dose is too low, hypothyroidism is undertreated. If the dose it too high, the thyroid hormone concentration is also too high, which leads to hyperactive thyroid effects [Ha00, Gr94].

There are two different tasks of determining an appropriate dose. The first one means to determine the initial dose, while later on the dose has to be updated continuously during a patient's lifetime. Precise determination of the initial dose is most important for newborn babies with congenital hypothyroidism, because for them every week of proper therapy counts.

#### **3.1 Computing an initial dose**

For the determination of an initial dose (fig.1.), a couple of prototypes, called guidelines, exist, which have been defined by commissions of experts. The assignment of a patient to a fitting guideline is obvious because of the way the guidelines have been defined. With the help of these guidelines a range for good doses can be calculated.

To compute an optimal dose, we retrieve similar cases with initial doses within the calculated ranges. Since there are only few attributes and since our case base is rather small, we use Tversky's sequential measure of dissimilarity [Tv77]. On the basis of those of the retrieved cases that had best therapy results an average initial therapy is calculated. Best therapy results can be determined by values of another blood test after two weeks of treatment with the initial dose. The opposite idea to consider cases with bad therapy results does not work here, because bad results may be caused by various reasons.

So, to compute an optimal dose recommendation, we apply two forms of adaptation. First, a calculation of ranges according to guidelines and patients attribute values. Secondly, we use Compositional Adaptation. That means, we take only similar cases with best therapy results into account and calculate the average dose for these cases, which has to be adapted to the query patient by another calculation.




Figure 1. Determination of an initial dose

**Example for computing an initial dose.** The query patient is a newborn baby that is 20 days old, has a weight of 4 kg and is diagnosed for hypothyroidism. The guideline for babies about 3 weeks age and normal weight recommend a levothyroxine therapy with a daily dose between 12 and 15  $\mu\text{g}/\text{kg}$ . So, because the baby weighs 4 kg, a range of 48-60  $\mu\text{g}$  would be calculated. The retrieval provides similar cases that must have doses within the calculated range. These cases are restricted to those where after two weeks treatment less than 10  $\mu\text{U}/\text{ml}$  thyroid stimulating hormone could be observed. Since these remaining similar cases are all treated alike, an average dose per kg is computed which subsequently is multiplied with the query's weight to deliver the optimal daily dose.

### 3.2 Updating the dose in a patient's lifetime

For monitoring the patient, three laboratory blood tests have to be made. Usually the results of these tests correspond to each other. Otherwise, it indicates a more complicated thyroid condition and additional tests are necessary. If the tests show that the patients thyroid hormone level is normal, it means that the current levothyroxine dose is OK. If the tests indicate that the thyroid hormone level is too low or too high, the current dose has to be increased resp. decreased by 25 or 50  $\mu\text{g}$  [Ha00, Gr94]. So, for monitoring, adaptation is just a simple calculation according to well-known standards.

### 3.3 Additional diseases or complications

It often occurs that patients do not only have hypothyroidism, but they suffer from further chronic diseases or complications. So, the levothyroxine therapy has to be checked for contraindications, adverse effects and interactions with additionally existing therapies. Since no alternative is available to replace levothyroxine, if necessary additionally existing therapies have to be modified, substituted, or compensated (figure 2) [Ha00, Gr94].

In our support program we perform three tests. The first one checks if another existing therapy is contraindicated to hypothyroidism. This holds only for very few therapies, namely for specific diets like soybean infant formula, but they are typical for newborn babies. Such diets have to be modified. Since no exact knowledge is available how to do it, our program just issues a warning saying that a modification is necessary.

The second test considers adverse effects. There are two ways to deal with them. A further existing therapy has either to be substituted or it has to be compensated by another drug. Since such knowledge is available, we have implemented corresponding rules for substitutional resp. compensational adaptation.

The third test checks for interactions between both therapies. Here we have implemented some adaptation rules, which mainly attempt to avoid the interactions. For example, if a patient has heartburn problems that are treated with an antacid, a rule for this situation exists that states that levothyroxine should be administered at least 4 hours after or before an antacid. However, if no adaptation rule can solve such an interaction problem, the same substitution rules as for adverse effects are applied.




Figure 2. Levothyroxine therapy and additionally existing therapies

## 4 Conclusion: Adaptation techniques for medical therapy problems

Our intention is to undertake first steps in the direction of developing a methodology for the adaptation problem for medical CBR systems. However, we have to admit that we are just at the beginning. Furthermore, our experience concerning the adaptation problem is mainly based on therapeutic tasks. Indeed, in contrast to ideas of many computer scientists, doctors are much more interested in therapeutic than in diagnostic support programs.

So, in this paper, we firstly reviewed how adaptation is handled in medical CBR systems and secondly enriched the experiences by additional examples from the endocrinology domain, where we have recently developed support programs. For medical therapy systems, that intend to apply the whole CBR cycle, at present we can summarise useful adaptation techniques. However, most of them are promising only for specific tasks.

**Abstraction** from single cases to more general prototypes seems to be a promising implicit support. However, if the prototypes correspond to guidelines, they may even explicitly solve some adaptation steps (see section 3.1).

**Compositional Adaptation** at first glance does not seem to be appropriate in medicine, because it was originally developed for configuration [WSC98]. However, it has been successfully applied for calculating therapy doses (e.g. in TA3-IVF [Ju98] and see section 3.1.).

**Constraints** are a promising adaptation technique too, but only for a specific situation (see section 2.2), namely for a set of solutions that can be reduced by checking e.g. contraindications (in ICONS) or contradictions (in GS.52).

**Adaptation rules.** The only technique that seems to be general enough to solve many medical adaptation problems is the application of adaptation rules or operators. Unfortunately, the technique is general, but the content of such rules has to be domain specific. Especially for complex medical tasks the generation of adaptation rules often is too time consuming and sometimes even impossible.

However, for therapeutic tasks some typical forms of adaptation rules can be made out, namely for substitutional and compensational adaptation (e.g. section 3.3.), and for calculating doses (e.g. section 3.2.). So, a next step might be an attempt to generate general adaptation operators for these typical forms of adaptation rules.

### References

- [AP94] Aamodt, A.; Plaza, E.: Case-based reasoning: Foundation issues. AICOM 7, 1994, S. 39-59.
- [Bi96] Bichindaritz I.: From cases to classes: Focusing on abstraction in case-based reasoning. In (Burkhard, H.-D.; Lenz, M. Hrsg.): Proc of 4th German Workshop on CBR, Humboldt University Berlin, 1996, S. 62-69.
- [Bi98] Bichindaritz, I. et al.: Case-based reasoning in CARE-PARTNER: Gathering evidence for evidence-based medical practice. In (Smyth, B.; Cunningham, P. Hrsg): Advances in Case-Based Reasoning, Proc of 4th European Workshop, Springer-Verlag, Berlin Heidelberg New York, 1998, S. 334-345.
- [Br90] Broder, A.: Strategies for efficient incremental nearest neighbor search. Pattern Recognition 23, 1990, S. 171-178.
- [BW98] Bergmann, R.; Wilke, W.: Towards a new formal model of transformational adaptation in case-based reasoning. In (Gierl, L.; Lenz, M. Hrsg.): Proc of 6th German Workshop on CBR, University of Rostock, 1998, S. 43-52.
- [FM99] Fuchs, B.; Mille, A.: A knowledge-level task model of adaptation in case-based

- reasoning. In (Althoff, K.-D. et al. Hrsg.): Case-Based Reasoning Research and Development, Proc of 3rd Int Conference, Springer-Verlag, Berlin Heidelberg New York, 1999, S. 118-131.
- [Du97] Dugas, M.: Clinical applications of Intranet-Technology. In (Dudeck, J. et al. Hrsg.): New Technologies in Hospital Information Systems. IOS Press, Amsterdam, 1997, S. 115-118.
- [GBS98] Gierl, L.; Bull, M; Schmidt, R.: CBR in Medicine. In (Lenz, M. et al. Hrsg.): Case-Based Reasoning Technology, From Foundations to Applications, Springer-Verlag, Berlin Heidelberg New York, 1998, S. 273-297.
- [Gr94] DeGroot, L.J.: Thyroid Physiology and Hypothyroidism. In (Besser, GM; Turner, M. Hrsg.): Clinical endocrinology. Wolfe, London, 1994, (Chapter 15).
- [GS94] Gierl, L.; Stengel-Rutkowski, S.: Integrating consultation and semi-automatic knowledge acquisition in a prototype-based architecture: Experiences with Dysmorphic Syndromes. Artificial Intelligence in Medicine 6, 1994, S. 29-49.
- [Ha00] Hampel, R.: Diagnostik und Therapie von Schilddrüsenfunktionsstörungen. UNI-MED Verlag, Bremen, 2000.
- [HK96] Hanney, K.; Keane, M.T.: Learning Adaptation Rules from a Case-Base. In (Smith, I.; Faltings, B. Hrsg.): Advances in Case-Based Reasoning, 3rd European Workshop, Springer-Verlag, Berlin Heidelberg New York, 1996, S. 179-192.
- [Ju98] Jurisica, I. et al: Case-based reasoning in IVF: prediction and knowledge mining. Artificial Intelligence in Medicine 12, 1998, S. 1-24
- [Ko88] Koton, P.: Reasoning about evidence in causal explanations. In (Kolodner, J. Hrsg.): First Workshop on CBR. Morgan Kaufmann, San Mateo, 1988, S. 260-270.
- [LB00] Lieber, J.; Bresson, B.: Case-based Reasoning for Breast Cancer Treatment Decision Help. In (Blanzieri, E.; Portinale, L. Hrsg.): Advances in Case-Based Reasoning, Proc of 5th European Workshop, Springer, Berlin, 2000, S. 173-185.
- [MM95] Macura, R.; Macura, K.: MacRad: Radiology image resource with a case-based retrieval system. In (Veloso, M.; Aamodt, A. Hrsg.): Case-Based Reasoning Research and Development, Proc of 1st Int Conference, Springer-Verlag, Berlin Heidelberg New York, 1995, S. 43-54.
- [Mo00] Montani, S. et al.: Diabetic patient's management exploiting Case-based Reasoning techniques. Computer Methods and Programs in Biomedicine 62, 2000, S. 205-218.
- [Pe01] Perner, P.: Why Case-Based Reasoning is Attractive for Image Interpretation. In (Aha, D.; Watson, I. Hrsg.): Case-Based Reasoning Research and Development, 4th Int Conference, Springer-Verlag, Berlin Heidelberg New York, 2001, S. 27-43.
- [PMS98] GJ Petot, G.J.; Marling, C; Sterling, L.: An artificial intelligence system for computer-assisted menu planing. Journal of American Diet Assoc 98 (9), 1998, S. 1009-10014
- [Qu93] Quinlan, J.: C4.5, Programs for Machine Learning. Morgan Kaufmann, San Mateo, 1993.
- [Sc99] Schmidt, R. et al.: Medical multiparametric time course prognoses applied to kidney function assessments. International Journal in Medical Informatics 53 (2-3), 1999, S. 253-264.
- [Sc01] Schmidt R. et al.: Case-Based Reasoning for Medical Knowledge-based Systems. International Journal of Medical Informatics 64 (2-3), 2001, S. 355-367
- [SG01] Schmidt, R.; Gierl, L.: Case-based Reasoning for Antibiotics Therapy Advice: An Investigation of Retrieval Algorithms and Prototypes. Artificial Intelligence in Medicine 23 (2), 2001, S. 171-186
- [SG02] Schmidt, R.; Gierl, L.: Prognostic Model for Early Warning of Threatening Influenza Waves. In (Minor, M.; Staab, S. Hrsg.): Proc 1st German Workshop on Experience Management, Köllen-Verlag, Bonn, 2002, S. 39-46.
- [SHK89] Stottler, R.H.; Henke, A.L.; King, J.A.: Rapid retrieval algorithms for case-based reasoning. In: Proc of 11th Int Joint Conference on Artificial Intelligence, Morgan



- Kaufmann, San Mateo, 1989, S. 233-237
- [SK98] Smyth, B.; Keane, M.T.: Adaptation-guided retrieval: questioning the similarity assumption in reasoning. *Artificial Intelligence* 102, 1998, S. 249-293
- [Tu88] R Turner: Organizing and using schematic knowledge for medical diagnosis. In (Kolodner, J. Hrsg.): *First Workshop on CBR*. Morgan Kaufmann, San Mateo, 1988, S. 435-446
- [Tv77] Tversky, A.: Features of similarity. *Psychological review* 84, 1977, S. 327-352
- [WSC98] Wilke, W.; Smyth, B.; Cunningham, P.: Using Configuration Techniques for Adaptation. In: M Lenz, B Bartsch-Spörl, H-D Burkhard, S Wess (eds.): *Case-Based Reasoning Technology, From Foundations to Applications*. Springer-Verlag, Berlin Heidelberg New York, 1998, S.139-168